

Level-1 Semester-2

# Pathology - MSS



*MCQ Lecture 6*  
**SOFT TISSUE TUMORS**

**DR M. YUSUF**



## MCQ on Soft Tissue Tumors

<p><b>1. <u>Soft tissues include:-</u></b></p> <ul style="list-style-type: none"> <li>a) Squamous epithelium</li> <li>b) Blood vessels</li> <li>c) Cartilage</li> <li>d) Brain tissue</li> <li>e) Glands</li> </ul>	<b>B</b>
<p><b>2. <u>Which of the followings is not a soft tissue tumor:-</u></b></p> <ul style="list-style-type: none"> <li>a) Liposarcoma</li> <li>b) Adenocarcinoma</li> <li>c) Rhabdomyosarcoma</li> <li>d) Leiomyoma</li> <li>e) Schwannoma</li> </ul>	<b>B</b>
<p><b>3. <u>Malignant skeletal muscle tumor is called:-</u></b></p> <ul style="list-style-type: none"> <li>a) Leiomyoma</li> <li>b) Leiomyosarcoma</li> <li>c) Schwannoma</li> <li>d) Angiosarcoma</li> <li>e) Rhabdomyosarcoma</li> </ul>	<b>E</b>
<p><b>4. <u>Benign tumor of peripheral nerves is called:-</u></b></p> <ul style="list-style-type: none"> <li>a) Lipoma</li> <li>b) Fibromatosis</li> <li>c) Leiomyoma</li> <li>d) Schwannoma</li> <li>e) Angioma</li> </ul>	<b>D</b>
<p><b>5. <u>Which of the following soft tissue tumors is intermediate grade lesion:-</u></b></p> <ul style="list-style-type: none"> <li>a) Liposarcoma</li> <li>b) Synovial sarcoma</li> <li>c) Fibromatosis</li> <li>d) Schwannoma</li> <li>e) Hemangioma</li> </ul>	<b>C</b>



<p>6. <u>Which of the following soft tissue tumors is of uncertain origin:-</u></p> <ul style="list-style-type: none"><li>a) Liposarcoma</li><li>b) Synovial sarcoma</li><li>c) Fibromatosis</li><li>d) Schwannoma</li><li>e) Hemangioma</li></ul>	E
<p>7. <u>Which of the followings is self-limited lesion requires minimal treatment:-</u></p> <ul style="list-style-type: none"><li>a) Rhabdomyosarcoma</li><li>b) Synovial sarcoma</li><li>c) Lipoma</li><li>d) Leiomyosarcoma</li><li>e) Angiosarcoma</li></ul>	C
<p>8. <u>Which of the following lesions has significant metastatic risk &amp; mortality:-</u></p> <ul style="list-style-type: none"><li>a) Lipoma</li><li>b) Leiomyoma</li><li>c) Schwannoma</li><li>d) Rhabdomyosarcoma</li><li>e) Fibromatosis</li></ul>	D
<p>9. <u>The most common soft tissue tumor in adult:-</u></p> <ul style="list-style-type: none"><li>a) Hemangioma</li><li>b) Schwannoma</li><li>c) Lipoma</li><li>d) Rhabdomyosarcoma</li><li>e) Synovial sarcoma</li></ul>	C
<p>10. <u>Lobulated soft mobile yellowish mass in SC tissue is consistent with:-</u></p> <ul style="list-style-type: none"><li>a) Leiomyoma</li><li>b) Lipoma</li><li>c) Fibromatosis</li><li>d) Schwannoma</li><li>e) Angiosarcoma</li></ul>	B



<p><b><u>Which of the followings is characterized by signet ring appearance under microscope:-</u></b></p> <p>a) Rhabdomyosarcoma b) Schwannoma c) Osteoma d) Lipoma e) Leiomyoma</p>	<b>D</b>
<p><b><u>12.S-100 protein is a marker for identification of:-</u></b></p> <p>a) Leiomyoma b) Leiomyosarcoma c) Rhabdomyosarcoma d) Lipoma e) Angiosarcoma</p>	<b>D</b>
<p><b><u>13.Recurring rapidly growing non-capsulated yellowish mass is most likely:-</u></b></p> <p>a) Lipoma b) Liposarcoma c) Leiomyosarcoma d) Schwannoma e) Hemangioma</p>	<b>B</b>
<p><b><u>14. The diagnostic cell in Liposarcoma is:-</u></b></p> <p>a) Lipocyte b) Liposome c) Lipoblast d) Mature fat cells e) UMC</p>	<b>C</b>
<p><b><u>15.Which of the followings is IHC marker for liposarcoma:-</u></b></p> <p>a) S-100 b) B-catenin c) SMA d) Myogenin e) CD31</p>	<b>A</b>



1. The

C

C

A

D

B

**16. Desmoid tumor refers to:-**

- a) Liposarcoma
- b) Superficial Fibromatosis
- c) Deep Fibromatosis
- d) Leiomyosarcoma
- e) Schwannoma

**17. Superficial Fibromatosis is most commonly seen in:-**

- a) Head
- b) Neck
- c) Palms
- d) Trunk
- e) Abdomen

**18. All of the followings are true about Fibromatosis except:-**

- a) Soft consistency
- b) Grey-white
- c) Single or multiple
- d) Locally aggressive
- e) Subcutaneous lesion

**19. Fascicles of bland fibroblasts surrounded by dense collagen bundles is characteristic of:-**

- a) Leiomyoma
- b) Leiomyosarcoma
- c) Rhabdomyosarcoma
- d) Fibromatosis
- e) Synovial sarcoma

**20. B-catenin is characteristic IHC marker for:-**

- a) Lipoma
- b) Fibromatosis
- c) Hemangioma
- d) Schwannoma
- e) Synovial sarcoma



<p><b><u>The most common soft tissue sarcoma of adolescent &amp; childhood is:-</u></b></p> <p>a) Liposarcoma b) Synovial sarcoma c) Rhabdomyosarcoma d) Angiosarcoma e) Leiomyosarcoma</p>	C
<p><b><u>22. Which of the following is not a microscopic type of Rhabdomyosarcoma:-</u></b></p> <p>a) Alveolar type b) Embryonic type c) Medullary type d) Pleomorphic type e) None of the above</p>	C
<p><b><u>23. Which of the following tumors has a microscopic appearance similar to lung tissue:-</u></b></p> <p>a) Leiomyosarcoma b) Pleomorphic Rhabdomyosarcoma c) Embryonal Rhabdomyosarcoma d) Alveolar Rhabdomyosarcoma e) Synovial sarcoma</p>	D
<p><b><u>24. Which of the followings is IHC marker for Rhabdomyosarcoma:-</u></b></p> <p>a) S-100 b) Desmin c) SMA d) FLi-1 e) CD99</p>	B
<p><b><u>25. Myogenin is a characteristic IHC marker for:-</u></b></p> <p>a) Leiomyosarcoma b) Liposarcoma c) Schwannoma d) Rhabdomyosarcoma e) Synovial sarcoma</p>	D



<p><b>26. <u>The most common site of leiomyoma is:-</u></b></p> <ul style="list-style-type: none"><li>a) Uterus</li><li>b) Skin</li><li>c) Stomach</li><li>d) Esophagus</li><li>e) Deep soft tissue</li></ul>	A
<p><b>27. <u>All of the followings are true about leiomyoma except:-</u></b></p> <ul style="list-style-type: none"><li>a) Grey-white</li><li>b) Bulging</li><li>c) Trabeculated</li><li>d) Hemorrhage &amp; necrosis</li><li>e) Firm consistency</li></ul>	D
<p><b>28. <u>Well-differentiated smooth muscle cells in fascicles without atypia is characteristic of:-</u></b></p> <ul style="list-style-type: none"><li>a) Leiomyosarcoma</li><li>b) Rhabdomyosarcoma</li><li>c) Leiomyoma</li><li>d) Schwannoma</li><li>e) Synovial sarcoma</li></ul>	C
<p><b>29. <u>Desmin is a characteristic IHC marker for:-</u></b></p> <ul style="list-style-type: none"><li>a) Leiomyoma</li><li>b) Lipoma</li><li>c) Schwannoma</li><li>d) Hemangioma</li><li>e) Synovial sarcoma</li></ul>	A
<p><b>30. <u>Which of the followings is not true for leiomyosarcoma:-</u></b></p> <ul style="list-style-type: none"><li>a) More common in children</li><li>b) Flemy mass</li><li>c) Grey-white</li><li>d) Hemorrhage and necrosis</li><li>e) Cellular pleomorphism</li></ul>	A



<p><b><u>What is the usual mitotic rate in leiomyosarcoma is:-</u></b></p> <p>a) 5 mitotic figures / 10 hpf  b) 1 mitotic figure / 10 hpf  c) 2 mitotic figures / 10 hpf  d) 4 mitotic figures / 10 hpf  e) 3 mitotic figures / 10 hpf</p>	A
<p><b><u>32. Which of the followings is IHC marker for leiomyosarcoma:-</u></b></p> <p>a) FLI-1  b) TLE1  c) CD31  d) SMA  e) S-100</p>	D
<p><b><u>33. Which of the followings is not true about schwannoma:-</u></b></p> <p>a) Benign tumor  b) Capsulated  c) Arise mainly in CNS  d) Elongated cells with wavy nuclei  e) Hyalinized vessels</p>	C
<p><b><u>34. Antoni A &amp; B areas are present in which of the following tumors:-</u></b></p> <p>a) Schwannoma  b) Lipoma  c) Leiomyoma  d) Hemangioma  e) Lymphangioma</p>	A
<p><b><u>35. Nuclear palisading around fibrillary process in schwannoma is known as:-</u></b></p> <p>a) Antoni A areas  b) Verocay bodies  c) Signet ring appearance  d) Antoni B areas  e) Desmoplasia</p>	B



<p><b>36. <u>Which of the following IHC markers is characteristic for schwannoma:-</u></b></p> <ul style="list-style-type: none"><li>a) SMA</li><li>b) S-100</li><li>c) CD31</li><li>d) CD34</li><li>e) CD99</li></ul>	B
<p><b>37. <u>The most common vascular tumor in infancy is:-</u></b></p> <ul style="list-style-type: none"><li>a) Angiosarcoma</li><li>b) Cavernous hemangioma</li><li>c) Lymphangioma</li><li>d) Capillary hemangioma</li><li>e) Angioblastoma</li></ul>	D
<p><b>38. <u>Strawberry lesion is a characteristic finding in:-</u></b></p> <ul style="list-style-type: none"><li>a) Lymphangioma</li><li>b) Schwannoma</li><li>c) Capillary hemangioma</li><li>d) Angiosarcoma</li><li>e) Cavernous hemangioma</li></ul>	C
<p><b>39. <u>Which of the followings is not true for hemangioma:-</u></b></p> <ul style="list-style-type: none"><li>a) Common vascular tumor</li><li>b) Common in head &amp; neck</li><li>c) Dark red color</li><li>d) Solid consistency</li><li>e) Blood filled spaces</li></ul>	D
<p><b>40. <u>Small vascular channels lined by flat endothelial cells is characteristic of:-</u></b></p> <ul style="list-style-type: none"><li>a) Capillary hemangioma</li><li>b) Cavernous hemangioma</li><li>c) Cavernous lymphangioma</li><li>d) Angiosarcoma</li><li>e) Synovial sarcoma</li></ul>	A



<p><u>Which of the following IHC markers is characteristic for capillary hemangioma:-</u></p> <p>a) S-100 b) CD31 c) CD99 d) SMA e) TLE1</p>	<b>B</b>
<p><b>42. <u>Port-wine nevus is a characteristic finding in:-</u></b></p> <p>a) Capillary hemangioma b) Lymphangioma c) Angiosarcoma d) Cavernous hemangioma e) Angioblastoma</p>	<b>D</b>
<p><b>43. <u>CD34 is a characteristic IHC marker for:-</u></b></p> <p>a) Lipoma b) Schwannoma c) Cavernous hemangioma d) Synovial sarcoma e) Leiomyosarcoma</p>	<b>C</b>
<p><b>44. <u>Large irregular spaces with proteinaceous intraluminal fluid containing lymphocytes is characteristic of:-</u></b></p> <p>a) Capillary hemangioma b) Synovial sarcoma c) Angiosarcoma d) Lymphangioma e) Cavernous hemangioma</p>	<b>D</b>
<p><b>45. <u>Highly aggressive malignancy of blood vessels is known as:-</u></b></p> <p>a) Angiosarcoma b) Rhabdomyosarcoma c) Liposarcoma d) Leiomyosarcoma e) Synovial sarcoma</p>	<b>A</b>



<p><b>46. <u>The presence of infiltrative vascular spaces with endothelial multilayering is characteristic of:-</u></b></p> <p>a) Lymphangioma b) Angiosarcoma c) Capillary hemangioma d) Cavernous hemangioma e) Synovial sarcoma</p>	B
<p><b>47. <u>Which of the followings is a characteristic IHC marker for Angiosarcoma:-</u></b></p> <p>a) S-100 b) SMA c) CD99 d) FLi-1 e) TLE1</p>	D
<p><b>48. <u>The most common site for synovial sarcoma is:-</u></b></p> <p>a) Head b) Neck c) Lower limb d) Upper limb e) Abdomen</p>	C
<p><b>49. <u>Which of the following IHC markers is characteristic for epithelial component in synovial sarcoma:-</u></b></p> <p>a) S-100 b) CD31 c) CD34 d) FLi-1 e) CK</p>	E
<p><b>50. <u>TLE1 is a characteristic IHC marker for:-</u></b></p> <p>a) Leiomyosarcoma b) Liposarcoma c) Synovial sarcoma d) Angiosarcoma e) Schwannoma</p>	C