

Motor Pathways

➔ **Tract:** Bundles or fasciculi of fibers have the same Origin, Termination & carry the same Function.

Pyramidal system (called so as they originate in pyramidal cells in cc) ⚡		Extrapyramidal tracts		Motor pathway to cerebellum																																																																																																														
	<table border="1"> <thead> <tr> <th></th> <th>Corticospinal</th> <th>Corticobulbar</th> </tr> </thead> <tbody> <tr> <td>WHICH MS??</td> <td>Muscles of body</td> <td>Muscles of head & neck</td> </tr> <tr> <td>Origin</td> <td></td> <td></td> </tr> <tr> <td>1ry motor area</td> <td>Area 4 (upper 2/3) → 40% ⚡</td> <td>Area 4 (Lower 1/3) → 40%</td> </tr> <tr> <td>Premotor area</td> <td colspan="2">Area 6 → 40%</td> </tr> <tr> <td>General sensory area</td> <td colspan="2">Area 3, 1 & 2 → 20%</td> </tr> <tr> <td>Course</td> <td colspan="2">Corona radiata of medullary center</td> </tr> <tr> <td>Internal capsule</td> <td>Posterior limb ⚡</td> <td>Genu</td> </tr> <tr> <td>Midbrain</td> <td colspan="2">Middle 3/5 of crus cerebri.</td> </tr> <tr> <td>Pons</td> <td colspan="2">Basis pontis</td> </tr> <tr> <td>Medulla</td> <td colspan="2">Pyramid</td> </tr> <tr> <td></td> <td>Motor decussation in lower medulla (85%)</td> <td>Ends in motor CN nuclei (3,4,5,6,7,9,10,11,12) - Not (1,2,8)</td> </tr> <tr> <td></td> <td></td> <td>▪ CN motor nuclei of 2 sides ⚡ mainly on opposite side except:</td> </tr> <tr> <td></td> <td> <table border="1"> <thead> <tr> <th>Opposite side</th> <th>Same side</th> </tr> </thead> <tbody> <tr> <td>85%</td> <td>15%</td> </tr> <tr> <td>Lateral</td> <td>Ventral</td> </tr> <tr> <td colspan="2">Corticospinal tract</td> </tr> </tbody> </table> </td> <td> <table border="1"> <thead> <tr> <th>7</th> <th>12</th> </tr> </thead> <tbody> <tr> <td>Lower part of fascial nucleus ↓ opposite side only</td> <td>Hypoglossal nucleus ↓ Genioglossus muscle</td> </tr> </tbody> </table> </td> </tr> <tr> <td>Termination</td> <td></td> <td></td> </tr> <tr> <td>Function</td> <td colspan="2"> <ul style="list-style-type: none"> 1 Facilitation of the flexor muscle tone 2 Initiation of highly skilled fine movements 3 Modulation of sensory input of the spinal cord </td> </tr> <tr> <td>Lesion</td> <td colspan="2"></td> </tr> <tr> <td>Cause</td> <td colspan="2"> <table border="1"> <tbody> <tr> <td>1 Trauma:</td> <td>- Fracture, stab or bullets</td> </tr> <tr> <td>2 Compression:</td> <td>- By tumor or herniated disc</td> </tr> <tr> <td>3 Vascular:</td> <td>- By occlusion of the artery (Most Common) ⚡</td> </tr> </tbody> </table> </td> </tr> <tr> <td>C/P ⚡</td> <td>- Above decussation → contralateral - Below decussation → ipsilateral</td> <td>- Weakness of the muscles supplied by CN. - UMN paralysis of the lower part of the contralateral facial nucleus, genioglossus</td> </tr> </tbody> </table>		Corticospinal	Corticobulbar	WHICH MS??	Muscles of body	Muscles of head & neck	Origin			1ry motor area	Area 4 (upper 2/3) → 40% ⚡	Area 4 (Lower 1/3) → 40%	Premotor area	Area 6 → 40%		General sensory area	Area 3, 1 & 2 → 20%		Course	Corona radiata of medullary center		Internal capsule	Posterior limb ⚡	Genu	Midbrain	Middle 3/5 of crus cerebri.		Pons	Basis pontis		Medulla	Pyramid			Motor decussation in lower medulla (85%)	Ends in motor CN nuclei (3,4,5,6,7,9,10,11,12) - Not (1,2,8)			▪ CN motor nuclei of 2 sides ⚡ mainly on opposite side except:		<table border="1"> <thead> <tr> <th>Opposite side</th> <th>Same side</th> </tr> </thead> <tbody> <tr> <td>85%</td> <td>15%</td> </tr> <tr> <td>Lateral</td> <td>Ventral</td> </tr> <tr> <td colspan="2">Corticospinal tract</td> </tr> </tbody> </table>	Opposite side	Same side	85%	15%	Lateral	Ventral	Corticospinal tract		<table border="1"> <thead> <tr> <th>7</th> <th>12</th> </tr> </thead> <tbody> <tr> <td>Lower part of fascial nucleus ↓ opposite side only</td> <td>Hypoglossal nucleus ↓ Genioglossus muscle</td> </tr> </tbody> </table>	7	12	Lower part of fascial nucleus ↓ opposite side only	Hypoglossal nucleus ↓ Genioglossus muscle	Termination			Function	<ul style="list-style-type: none"> 1 Facilitation of the flexor muscle tone 2 Initiation of highly skilled fine movements 3 Modulation of sensory input of the spinal cord 		Lesion			Cause	<table border="1"> <tbody> <tr> <td>1 Trauma:</td> <td>- Fracture, stab or bullets</td> </tr> <tr> <td>2 Compression:</td> <td>- By tumor or herniated disc</td> </tr> <tr> <td>3 Vascular:</td> <td>- By occlusion of the artery (Most Common) ⚡</td> </tr> </tbody> </table>		1 Trauma:	- Fracture, stab or bullets	2 Compression:	- By tumor or herniated disc	3 Vascular:	- By occlusion of the artery (Most Common) ⚡	C/P ⚡	- Above decussation → contralateral - Below decussation → ipsilateral	- Weakness of the muscles supplied by CN. - UMN paralysis of the lower part of the contralateral facial nucleus, genioglossus	<ul style="list-style-type: none"> 1 Facilitate the flexor muscle tone: <ul style="list-style-type: none"> - Rubrospinal - Tectospinal - Medullary vestibulospinal “Sulco-marginal” - Medullary reticulospinal “lateral” 2 Facilitate the extensor muscle tone: <ul style="list-style-type: none"> - Lateral vestibulospinal - Pontine (medial) reticulospinal <table border="1"> <thead> <tr> <th></th> <th>Origin</th> <th>Function</th> </tr> </thead> <tbody> <tr> <td>Tectospinal</td> <td>Superior colliculus of opposite side.</td> <td>Reflexes ass with vision & hearing</td> </tr> <tr> <td>Rubrospinal</td> <td>Red nucleus of opposite side. ⚡</td> <td>Carries information from CC & cerebellum to spinal cord</td> </tr> <tr> <td>Medullary reticulospinal</td> <td>Medullary reticular formation</td> <td>Facilitate flexor muscles tone</td> </tr> <tr> <td>Medullary vesibulospinal</td> <td>Medial vestibular nucleus.</td> <td></td> </tr> <tr> <td>Pontine vesibulospinal</td> <td>Lateral vestibular nucleus.</td> <td>Facilitate extensor muscles tone</td> </tr> <tr> <td>Pontine reticulospinal</td> <td>Pontine reticular formation.</td> <td></td> </tr> </tbody> </table>		Origin	Function	Tectospinal	Superior colliculus of opposite side.	Reflexes ass with vision & hearing	Rubrospinal	Red nucleus of opposite side. ⚡	Carries information from CC & cerebellum to spinal cord	Medullary reticulospinal	Medullary reticular formation	Facilitate flexor muscles tone	Medullary vesibulospinal	Medial vestibular nucleus.		Pontine vesibulospinal	Lateral vestibular nucleus.	Facilitate extensor muscles tone	Pontine reticulospinal	Pontine reticular formation.		<ul style="list-style-type: none"> ☐ Cortico-Ponto-Cerebellar Pathway: <table border="1"> <thead> <tr> <th></th> <th>From</th> <th>Transmit</th> </tr> </thead> <tbody> <tr> <td>Fronto-pontine fibers</td> <td>Frontal lobe.</td> <td>Motor information</td> </tr> <tr> <td>Non fronto-pontine fibers</td> <td>Parietal, temporal & occipital lobes.</td> <td>Sensory information</td> </tr> </tbody> </table> <p>Descending Autonomic Fibers</p> <table border="1"> <tbody> <tr> <td>Origin</td> <td>- From hypothalamus and autonomic centers in reticular formation.</td> </tr> <tr> <td>Termination</td> <td>- Sympathetic nucleus in lateral horn of spinal cord (T1:L3). - Sacral parasympathetic nucleus (S2:S4).</td> </tr> <tr> <td>Lesion</td> <td>- Above T1 will cause Horner's Syndrome: ⚡ 1) Ptosis: paralysis of muller's muscle. 2) Miosis: constriction of pupil. 3) Anhidrosis: dry face due to paralysis of sweat glands. 4) Enophthalmos: retraction of the eyeball. 5) Flushing: warm and red face</td> </tr> </tbody> </table>		From	Transmit	Fronto-pontine fibers	Frontal lobe.	Motor information	Non fronto-pontine fibers	Parietal, temporal & occipital lobes.	Sensory information	Origin	- From hypothalamus and autonomic centers in reticular formation.	Termination	- Sympathetic nucleus in lateral horn of spinal cord (T1:L3). - Sacral parasympathetic nucleus (S2:S4).	Lesion	- Above T1 will cause Horner's Syndrome: ⚡ 1) Ptosis: paralysis of muller's muscle. 2) Miosis: constriction of pupil. 3) Anhidrosis: dry face due to paralysis of sweat glands. 4) Enophthalmos: retraction of the eyeball. 5) Flushing: warm and red face
	Corticospinal	Corticobulbar																																																																																																																
WHICH MS??	Muscles of body	Muscles of head & neck																																																																																																																
Origin																																																																																																																		
1ry motor area	Area 4 (upper 2/3) → 40% ⚡	Area 4 (Lower 1/3) → 40%																																																																																																																
Premotor area	Area 6 → 40%																																																																																																																	
General sensory area	Area 3, 1 & 2 → 20%																																																																																																																	
Course	Corona radiata of medullary center																																																																																																																	
Internal capsule	Posterior limb ⚡	Genu																																																																																																																
Midbrain	Middle 3/5 of crus cerebri.																																																																																																																	
Pons	Basis pontis																																																																																																																	
Medulla	Pyramid																																																																																																																	
	Motor decussation in lower medulla (85%)	Ends in motor CN nuclei (3,4,5,6,7,9,10,11,12) - Not (1,2,8)																																																																																																																
		▪ CN motor nuclei of 2 sides ⚡ mainly on opposite side except:																																																																																																																
	<table border="1"> <thead> <tr> <th>Opposite side</th> <th>Same side</th> </tr> </thead> <tbody> <tr> <td>85%</td> <td>15%</td> </tr> <tr> <td>Lateral</td> <td>Ventral</td> </tr> <tr> <td colspan="2">Corticospinal tract</td> </tr> </tbody> </table>	Opposite side	Same side	85%	15%	Lateral	Ventral	Corticospinal tract		<table border="1"> <thead> <tr> <th>7</th> <th>12</th> </tr> </thead> <tbody> <tr> <td>Lower part of fascial nucleus ↓ opposite side only</td> <td>Hypoglossal nucleus ↓ Genioglossus muscle</td> </tr> </tbody> </table>	7	12	Lower part of fascial nucleus ↓ opposite side only	Hypoglossal nucleus ↓ Genioglossus muscle																																																																																																				
Opposite side	Same side																																																																																																																	
85%	15%																																																																																																																	
Lateral	Ventral																																																																																																																	
Corticospinal tract																																																																																																																		
7	12																																																																																																																	
Lower part of fascial nucleus ↓ opposite side only	Hypoglossal nucleus ↓ Genioglossus muscle																																																																																																																	
Termination																																																																																																																		
Function	<ul style="list-style-type: none"> 1 Facilitation of the flexor muscle tone 2 Initiation of highly skilled fine movements 3 Modulation of sensory input of the spinal cord 																																																																																																																	
Lesion																																																																																																																		
Cause	<table border="1"> <tbody> <tr> <td>1 Trauma:</td> <td>- Fracture, stab or bullets</td> </tr> <tr> <td>2 Compression:</td> <td>- By tumor or herniated disc</td> </tr> <tr> <td>3 Vascular:</td> <td>- By occlusion of the artery (Most Common) ⚡</td> </tr> </tbody> </table>		1 Trauma:	- Fracture, stab or bullets	2 Compression:	- By tumor or herniated disc	3 Vascular:	- By occlusion of the artery (Most Common) ⚡																																																																																																										
1 Trauma:	- Fracture, stab or bullets																																																																																																																	
2 Compression:	- By tumor or herniated disc																																																																																																																	
3 Vascular:	- By occlusion of the artery (Most Common) ⚡																																																																																																																	
C/P ⚡	- Above decussation → contralateral - Below decussation → ipsilateral	- Weakness of the muscles supplied by CN. - UMN paralysis of the lower part of the contralateral facial nucleus, genioglossus																																																																																																																
	Origin	Function																																																																																																																
Tectospinal	Superior colliculus of opposite side.	Reflexes ass with vision & hearing																																																																																																																
Rubrospinal	Red nucleus of opposite side. ⚡	Carries information from CC & cerebellum to spinal cord																																																																																																																
Medullary reticulospinal	Medullary reticular formation	Facilitate flexor muscles tone																																																																																																																
Medullary vesibulospinal	Medial vestibular nucleus.																																																																																																																	
Pontine vesibulospinal	Lateral vestibular nucleus.	Facilitate extensor muscles tone																																																																																																																
Pontine reticulospinal	Pontine reticular formation.																																																																																																																	
	From	Transmit																																																																																																																
Fronto-pontine fibers	Frontal lobe.	Motor information																																																																																																																
Non fronto-pontine fibers	Parietal, temporal & occipital lobes.	Sensory information																																																																																																																
Origin	- From hypothalamus and autonomic centers in reticular formation.																																																																																																																	
Termination	- Sympathetic nucleus in lateral horn of spinal cord (T1:L3). - Sacral parasympathetic nucleus (S2:S4).																																																																																																																	
Lesion	- Above T1 will cause Horner's Syndrome: ⚡ 1) Ptosis: paralysis of muller's muscle. 2) Miosis: constriction of pupil. 3) Anhidrosis: dry face due to paralysis of sweat glands. 4) Enophthalmos: retraction of the eyeball. 5) Flushing: warm and red face																																																																																																																	